

Parameterization of copula functions for bivariate survival data in the surrosurv package

Federico Rotolo

November 9, 2016

Let define the joint survival function of S and T via a copula function:

$$S(s, t) = P(S \geq s, T \geq t) = C(u, v)|_{u=S_S(s), v=S_T(t)}, \quad (1)$$

where $S_S(\cdot) = P(S \geq s)$ and $S_T(\cdot) = P(T \geq t)$ are the marginal survival functions of S and T .

In the case of possibly right-censored data, the individual contribution to the likelihood is

- $S(s, t) = C(u, v)|_{S_S(s), S_T(t)}$ if S is censored at time s and T is censored at time t ,
- $-\frac{\partial}{\partial t} S(s, t) = \frac{\partial}{\partial v} C(u, v)|_{S_S(s), S_T(t)} f_S(s)$ if S is censored at time s and $T = t$,
- $-\frac{\partial}{\partial s} S(s, t) = \frac{\partial}{\partial u} C(u, v)|_{S_S(s), S_T(t)} f_T(t)$ if $S = s$ and T is censored at time t ,
- $\frac{\partial^2}{\partial s \partial t} S(s, t) = \frac{\partial^2}{\partial u \partial v} C(u, v)|_{S_S(s), S_T(t)} f_S(s) f_T(t)$ if $S = s$ and $T = t$.

Clayton copula

The bivariate Clayton [1978] copula is defined as

$$C(u, v) = (u^{-\theta} + v^{-\theta} - 1)^{-1/\theta}, \quad \theta > 0. \quad (2)$$

The first derivative with respect to u is

$$\begin{aligned} \frac{\partial}{\partial u} C(u, v) &= (u^{-\theta} + v^{-\theta} - 1)^{-\frac{1+\theta}{\theta}} u^{-(1+\theta)} \\ &= \left[\frac{C(u, v)}{u} \right]^{1+\theta}. \end{aligned} \quad (3)$$

The second derivative with respect to u and v is

$$\frac{\partial^2}{\partial u \partial v} C(u, v) = (1 + \theta) \frac{C(u, v)^{1+2\theta}}{(uv)^{1+\theta}}. \quad (4)$$

The Kendall [1938]'s tau for the Clayton copula is

$$\tau = \frac{\theta}{\theta + 2}. \quad (5)$$

Plackett copula

The bivariate Plackett [1965] copula is defined as

$$C(u, v) = \frac{[Q - R^{1/2}]}{2(\theta - 1)}, \quad \theta > 0, \quad (6)$$

with

$$\begin{aligned} Q &= 1 + (\theta - 1)(u + v), \\ R &= Q^2 - 4\theta(\theta - 1)uv. \end{aligned} \quad (7)$$

Given that

$$\frac{\partial}{\partial u}Q = \theta - 1, \quad (8)$$

$$\frac{\partial}{\partial u}R = 2(\theta - 1)(1 - (\theta + 1)v + (\theta - 1)u), \quad (9)$$

the first derivative of $C(u, v)$ with respect to u is

$$\begin{aligned} \frac{\partial}{\partial u}C(u, v) &= \frac{1}{2} \left[1 - \frac{1 - (\theta + 1)v + (\theta - 1)u}{R^{1/2}} \right] \\ &= \frac{1}{2} \left[1 - \frac{Q - 2\theta v}{R^{1/2}} \right]. \end{aligned} \quad (10)$$

By defining

$$f = 1 - (\theta + 1)v + (\theta - 1)u, \quad (11)$$

$$g = R^{1/2} \quad (12)$$

and given that

$$f' = \frac{\partial}{\partial u}f = -(\theta + 1), \quad (13)$$

$$g' = \frac{\partial}{\partial u}g = \frac{\theta - 1}{R^{1/2}}(1 - (\theta + 1)u + (\theta - 1)v), \quad (14)$$

then, the second derivative with respect to u and v is

$$\begin{aligned} \frac{\partial^2}{\partial u \partial v}C(u, v) &= -\frac{f'g - fg'}{2g^2} \\ &= -\frac{\theta}{R^{3/2}}[1 + (\theta - 1)(u + v - 2uv)] \\ &= -\frac{\theta}{R^{3/2}}[Q - 2(\theta - 1)uv]. \end{aligned} \quad (15)$$

The Kendall's tau for the Plackett copula cannot be computed analytically and is obtained numerically.

Gumbel–Hougaard copula

The bivariate Gumbel [1960]–Hougaard [1986] copula is defined as

$$C(u, v) = \exp(-Q^\theta), \quad \theta \in (0, 1), \quad (16)$$

with

$$Q = (-\ln u)^{1/\theta} + (-\ln v)^{1/\theta}. \quad (17)$$

Given that

$$\frac{\partial}{\partial u}Q = -\frac{(-\ln u)^{1/\theta-1}}{\theta u}, \quad (18)$$

then, the first derivative with respect to u is

$$\frac{\partial}{\partial u}C(u, v) = \frac{(-\ln u)^{1/\theta-1}}{u}C(u, v)Q^{\theta-1} \quad (19)$$

and the second derivative with respect to u and v is

$$\frac{\partial^2}{\partial u \partial v}C(u, v) = \frac{[(-\ln u)(-\ln v)]^{1/\theta-1}}{uv}C(u, v)Q^{\theta-2}\left[\frac{1}{\theta} - 1 + Q^\theta\right]. \quad (20)$$

The Kendall's tau for the Gumbel–Hougaard copula is

$$\tau = 1 - \theta. \quad (21)$$

References

- DG Clayton. A model for association in bivariate life tables and its application in epidemiological studies of familial tendency in chronic disease incidence. *Biometrika*, 65:141–151, 1978. doi:10.1093/biomet/65.1.141.
- EJ Gumbel. Distributions des valeurs extrêmes en plusieurs dimensions. *Publ Inst Statist Univ Paris*, 9:171–173, 1960.
- P Hougaard. A class of multivariate failure time distributions. *Biometrika*, 73:671–678, 1986. doi:10.1093/biomet/73.3.671.
- MG Kendall. A new measure of rank correlation. *Biometrika*, 30(1/2):81–93, 1938. URL <http://www.jstor.org/stable/2332226>.
- R. B. Nelsen. *An Introduction to Copulas*. Springer Verlag, second edition, 2006. doi:10.1007/0-387-28678-0.
- RL Plackett. A class of bivariate distributions. *J Am Statist Assoc*, 60:516–522, 1965. doi:10.1080/01621459.1965.10480807.